

PATENT APPLICATION

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Docket No: Q96434

Masaaki HIRANO, et al.

Appln. No.: 10/588,485

Group Art Unit: 1614

Confirmation No.: 8206

Examiner: not yet assigned

Filed: August 4, 2006

For:

PROPANE-1, 3-DIONE DERIVATIVE OR SALT THEREOF

INFORMATION DISCLOSURE STATEMENT UNDER 37 C.F.R. §§ 1.97 and 1.98

MAIL STOP AMENDMENT

Commissioner for Patents P.O. Box 1450 Alexandria, VA 22313-1450

Sir:

In accordance with the duty of disclosure under 37 C.F.R. § 1.56, Applicants hereby notify the U.S. Patent and Trademark Office of the documents which are listed on the attached PTO/SB/08 A & B (modified) form and/or listed herein and which the Examiner may deem material to patentability of the claims of the above-identified application.

One copy of each of the listed documents is submitted herewith, except for the following: U.S. patents and/or U.S. patent publications; and co-pending non-provisional U.S. applications filed after June 30, 2003.

The present Information Disclosure Statement is being filed: (1) No later than three months from the application's filing date; (2) Before the mailing date of the first Office Action on the merits (whichever is later); or (3) Before the mailing date of the first Office Action after

Attorney Docket No.: Q96434 INFORMATION DISCLOSURE STATEMENT

U.S. Appln. No.: 10/588,485

filing a request for continued examination (RCE) under §1.114, and therefore, no Statement

under 37 C.F.R. § 1.97(e) or fee under 37 C.F.R. § 1.17(p) is required.

In compliance with the concise explanation requirement under 37 C.F.R. § 1.98(a)(3) for

foreign language documents, Applicants submit the following explanations:

English language abstracts submitted herewith, constitute a concise explanation for the

foreign language documents on the attached list.

The submission of the listed documents is not intended as an admission that any such

document constitutes prior art against the claims of the present application. Applicants do not

waive any right to take any action that would be appropriate to antedate or otherwise remove any

listed document as a competent reference against the claims of the present application.

The USPTO is directed and authorized to charge all required fees, except for the Issue

Fee and the Publication Fee, to Deposit Account No. 19-4880. Please also credit any

overpayments to said Deposit Account.

Respectfully submitted,

Registration No. 32,197

Mark Boland

SUGHRUE MION, PLLC

Telephone: (202) 293-7060

Facsimile: (202) 293-7860

WASHINGTON DC SUGHRUE/265550

65565

CUSTOMER NUMBER

Date: August 24, 2007

•	
	Non-English references
	WO98/04562
	DD224422
	JP2000-95767
	JP09061992
	JP1992-334369
	JP1989-205130
	JP2002241758
	JP63271341
	JP56161538
	JP2002088284
	JP2004061583
	JP2002268239
	JP03259150
	JP03164722
1	JP02079007
	JP02189547
	JP02054268
	JP63032542
	JP62006254
٠	ID50064840

Statement By APPLICANT

(use as many sheets as necessary)

AUG 2 4 2007

of

Sheet

Complete if Known					
Application Number	10/588,485				
Confirmation Number	8206				
Filing Date	August 4, 2006				
Pirst Named Inventor	Masaaki HIRANO				
	1614				
Attorney Docket Number	not yet assigned				
Attorney Docket Number	Q96434				

	U.S. PATENT DOCUMENTS						
Examiner Initials*	Cite No.1	Document N Number	Kind Code ² (if known)	Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document		
		US 20030191164	A1	10-09-2003	Msaaki HIRANO et al.		
US 4062686 12-1		12-13-1977	Eastman Kodak Company				
		US 4119466		10-10-1978	Eastman Kodak Company		
		US 5202221	A	04-13-1993	Fuji Photo Film Co., Ltd.		
		US 5616537	A	04-01-1997	Kumiai Chemical Industry Co., Ltd., et al.		
		US 5770544	A	06-23-1998	Kumiai Chemical Industry Co., Ltd., et al.		
		US 5817819	A	10-06-1998	Takeda Chemical Industries, Ltd.,		
		US 5141841	A	08-25-1992	Vickers PLC		
		US 5519136	A	05-21-1996	John R. Wade		
		US 5055579		10-08-1991	Hoechst Aktiengesellschaft		
		US 5593818	Α	01-14-1997	Fuji Photo Film Co., Ltd		
		US 6087503	A	07-11-2000	Takeda Chemical Industries Ltd.		
	•	US 6162813	A	12-19-2000	Merck & Co., Inc		
		US 6413503	B1	07-02-2002	BASF Aktiengesellschaft		
US 6153371 A 11		11-28-2000	Eastman Kodak Company				
		11-30-1999	Eastman Kodak Company				
		US 5747236	Α	05-05-1998	Eastman Kodak Company		
		US 5747235	Α	05-05-1998	Eastman Kodak Company		
		US 6051359	Α	04-18-2000	Fuji Photo Film Co., Ltd.		
		US 5593818	Α	01-14-1997	Fuji Photo Film Co.,. Ltd.		
		US 5385807	Α	01-31-1995	Fuji Photo Film Co.,. Ltd.		
Ī		US 4950640		08-21-1990	Eastman Kodak Company		
		US 6140384	A	10-31-2000	Fuji Photo Film Co.,. Ltd.		
		US 5445930	A	08-29-1995	Fuji Photo Film Co.,. Ltd.		
		US 5738982	A	04-14-1998	Fuji Photo Film Co.,. Ltd.		
		US 6346534	B1	02-12-2002	Neurocrine Biosciences, Inc		
		US 6395733	B1	05-28-2002	Pfizer Inc.		
		US 20020177556	A1	11-28-2002	Jurgen Engel et al.		
		US 20040029040	A1	02-12-2004	FUJI PHOTO FILM CO., LTD		
		US 5104783		04-14-1992	Fuji Photo Film Co.,. Ltd.		
		US 4636509		01-13-1987	Glaxo Group Limited		
		US 4263393		04-21-1981	Eastman Kodak Company		
		US 6413503	Bl	07-02-2002	BASF Aktiengesellschaft		
		US 6468711	B1	10-22-2002	Fuji Photo Film Co.,. Ltd.		
		US 5064747		11-12-1991	Fuji Photo Film Co.,. Ltd.		
		US 5112743		05-12-1992	Fuji Photo Film Co.,. Ltd.		
		US 4966828		10-30-1990	Fuji Photo Film Co.,. Ltd.		
		US 4946960		08-07-1990	Vickers PLC		

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

¹Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial number of the patent document. Kind of document by the appropriate symbols as indicated to the following the f

	FOREIGN PATENT DOCUMENTS							
Examiner Initials*	Cite No.1		Country	reign Patent Docur	Kind Code ⁵	Publication Date MM-DD-YYYY	Name of Patentee or Applicant of Cited Document	Translation ⁶
		Code ³		(if known)		DIVI DIVOTO DI LA COLUMN	<u> </u>	
		EP	333156	A2	09-20-1989	FUJI PHOTO FILM CO., LTD	abstract	
		WO	9952888	A1	10-21-1999	AYUKO	abstract	
		EP	780730	A2	06-25-1997	FUJI PHOTO FILM CO., LTD	Abstract	
		WO	2005097090	A2	10-20-2005	ICOS CORPORATION	Abstract	
		EP	713143	A2	05-22-1996	FUJI PHOTO FILM CO., LTD.	Abstract	
		wo	9804562	A1	02-05-1998	BAYER AG	Abstract	
		DD	224422	A1	03-07-1985	VEB FILMFABRIK WOLFEN	Abstract	
		JP	2000-095767	Α	04-04-2000	TAKEDA CHEM IND LTD	Abstract	
		JP	09-061992	Α	03-07-1997	FUJI PHOTO FILM CO., LTD.	Abstract	
		JP	04-334369	Α	11-20-1992	FUJI PHOTO FILM CO., LTD.	Abstract	
		JP	01-205130	A	08-17-1989	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	2002-241758	A	08-28-2002	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	63-271341	Α	11-09-1988	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	56-161538	Α	12-11-1981	MITSUBISHI CHEM IND LTD	Abstract	
		JP	2002-088284	A	03-27-2002	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	2004-061583	Α	02-26-2004	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	2002-268239	Α	09-18-2002	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	03-259150	Α	11-19-1991	MITSUBISHI KASEI CORP	Abstract	
		JP	02-079007	Α	03-19-1990	SUMITOMO ELECTRIC IND LTD	Abstract	
		JP	02-189547	Α	07-25-1990	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	02-054268	Α	02-23-1990	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	03-164722	Α	07-16-1991	FUJI PHOTO FILM CO., LTD	Abstract	
		JP	63-032542	A	02-12-1988	MITSUBISHI CHEM IND LTD	Abstract	
		JP	62-006254	Α	01-13-1987	MITSUBISHI CHEM IND LTD	Abstract	
		JP	59-064840	Α	04-12-1984	MITSUBISHI CHEM IND LTD	Abstract	

	NON PATENT LITERATURE DOCUMENTS					
Examiner Initials*	Cite No. ¹	Include name of the author (in CAPITAL LETTERS), title of the article (when appropriate), title of the item (book, magazine, journal, serial, symposium, catalog, etc.), date, page(s), volume-issue number(s), publisher, city, and/or country where published.	Translation ⁶			
		Simone, Oncology: Introduction, Cecil Textbook of Medicine, 20.sup.th Edition, vol. 1, pp. 1004-1010, 1996				
		Huirne et al., PubMed Abstract (Lancet 358(9295):1793-803), Nov. 2001 Gonadotropin-releasing-				
	hormone-receptor antagonists					
		Junko Ishida et al., "Antitumor Agents. Part 214: .supdagger. Synthesis and Evaluation of Curcumin Analogues as Cytotoxic Agents" Bioorganic & Medicinal Chemistry 10 (2002) 3481-3487.				
		Chemistry of Heterocyclic Compounds (New York, NY, United States)(Translation of Khimiya Geterotsiklicheskikh Soedinenii) (2001), 37(5), 554-559				
		1.B. Dzvinchuk et al., "Formation of Unsymmetrical 2-(Diacylmethylene)-2,3-Dihydro-1H-Benzimidazoles During Acidolysis of 1-Benzoyl2-(.betaBenzoyloxybetaPhenylvinyl)-1H-Benzimidazole",				
		Jaro Komenda et al., Electrochemical Behavior and ESR Spectra of Nitro Substituted Mono-to and Debenzoylmethylenebenzthiazolines and Selenazolies, Collect. Czech. Chem. Commun. (1979), vol. 44(5),				
		pp. 1540-51.				

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the Emperor must precede the serial interpretation of the year of the reign of the emperor must precede the serial interpretation of the year of the reign of the year of the year of the year of the reign of the year of the year

Collect. Czech. Chem. Commun. (1973), vol. 38(12), pp. 3616-3622. . A. Mistr et al., Organische Lichtempfindliche Stoffe V. Aclmethylenderivate Heterocyclischer Stickstoffhaltiger Basen ALS Sensibilisatoren Lichtempfindlicher Polymerer, Organic light-sensitive substances. Acylmethylene derivatives of heterocyclic nitrogen-containing bases as sensitizers for light-sensitive polymers. ABS Condensation of quaternary salts of 2-methylbenzothiazoles or selenazoles with substituted benzyl chlorides in pyridine gave 18 mono- and diacylmethylene derivs. of the general formula I (Y = S, Se; R1 = Me, Et; R2 = H, p-OMe, m-NO2, p-NO2, p-I; R3 = H, R2C6H4(SH:CH)nCO; n = 0,1), which were examd as sensitizing agents for light -sensitive poly(vinyl cinnamate) (II) [24968-99-8] and poly(vinyl pazidobenzoate) [29928-09-4]. A dropwise addition of 0.033 mole quaternary benzothiazole salt to 0.033 mole corresponding acyl chloride at 10.deg. followed by 1 hr strring at 20.deg. gave 49.9% 2-[(4methoxybenzoyl)methylene]-3-ethylbenzothiazoline (I; Y = S; R1 = Et; R2 = p-OCH3; R3 = H; n = o) [51936-64-2]. The light sensitivity data obtained for II indicated that Se, present in the heterocyclic I ring, was more effective than S, and that the substitution on the benzoyl ring decreased the light sensitivity effect of I through the substituent series p-OMe, m-NO2, CH:CH, p-I, p-NO2. A. Mistr et al., Organische Lichtempfindliche Stoffee II. Benzoylmethylenderivate Heretocyclischer Stickstoffhaltiger Basen ALS Sensibilisatoren Fur Lichtempfinliche Polymere, Collect. Czech. Commun. (1971), vol. 36(1), pp. 150-163. **ABS** Alkyl toluate salts of substituted and unsubstituted benzothiazole and benzoselenazole are treated with BzCl in pyridine to prepare I (Y = S or Se, R = H or Bz, R1 = Me or Et) and analogs containing a Cl, Me, Et, MeO, or benzo group on the 6-membered ring. I (R = Bz) had greater light sensitizing activity in poly(vinyl cinnamate) and poly(vinyl p-azidobenzoate) than did I (R = H), and the benzoselenazoline photosensitizers were more active than the benzothiazoline photosensitizers. Substituents at the 5-position on the aromatic ring had no effect and 6-methoxy substituents, a small pos. effect on sensitizing activity. Varying the R1 alkyl substituent affected solubility but not sensitizing activity.

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant

Applicant's unique citation designation number (optional). 2See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. 3 Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). 4For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the kerial to the north the first t

G.I. Gaeva and K.S. Liadikov, Zh. Nauch. Prikl. Fotogr. Kinematogr. (1971) vol. 16(4), pp. 282-288 Sensitization of poly(vinyl cinnamate) by derivatives of benzoyl- and dibenzoylmethylenebenzothiazoline and -benzoselenazoline. ABS Of the 22 benzothiazoline dyes and benzoselenazoline dyes studied, of the general formula (I) (where Y = S or Se, R = Me or Et, R1 = H or COPh, R2 = COPh, R3 = H, Cl, MeO, and 4,5- or 6,7-benzo group), 1-methyl-2- (dibenzoylmethylenenaphtho[1,2-d]thiazole and 3-methyl-5-methoxy-2- (dibenzoylmethylene)benzoselenazole had the highest sensitization effectiveness and increased the light sensitivity of poly(vinyl cinnamate) (I) 2.5 times. The spectral sensitivity of dyes and their optimum concentration in I were determined A sensitization mechanism was proposed.
The Chemistry and Biological Activity of Synthetic and Natural Compounds: Nitrogen-Containing Heterocycles, Vol. 1 (2006), pp. 243-248.
Chemistry of Heterocyclic Compounds (New York, NY, United States)(Translation of Khimiya Geterotsiklicheskikh Soedinenii) (2001), 37(5), 554-559
Bioorganic & Medicinal Chemistry Letters, Volume 15, Issue 11, 2 June 2005, Pages 2894-2897 Synthesis, in vivo and in vitro biological activity of novel azaline B analogs

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Zh. Organic Khim. (1994), 30(6), 909-14

LA Russian

Explanation: Found by CAS Search. CAS Search Result is set forth below.

L13 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 2006 ACS on STN

AN 1995:373237 HCAPLUS Full-text

DN 123:169551

ED Entered STN: 24 Feb 1995

TI C-Monobenzoylation and dibenzoylation of 2-methylbenzimidazole by benzoyl chloride

AU Dzvinchuk, I. B.; Lozinskii, M. O.; Vypirailenko, A. V.

CS Inst. Org. Khim., Kiev, Ukraine

SO Zhurnal Organicheskoi Khimii (1994), 30(6), 909-14 CODEN: ZORKAE; ISSN: 0514-7492

PB Nauka

DT Journal

LA Russian

CC 28-9 (Heterocyclic Compounds (More Than One Hetero Atom))

GI

- AB Reaction of 2-methylbenzimidazole with BzCl in the presence of Et3N gave monobenzoyl (I), dibenzoyl (II and III), tribenzoyl (IV), and tetrabenzoyl derivs. (V). The interconversion of these products and the effect of temperature were examined
- ST benzoylation methylbenzimidazole; benzimidazole methyl benzoylation

IT Benzoylation

(of methylbenzimidazole by benzoyl chloride)

IT 615-15-6, 2-Methylbenzimidazole

RL: RCT (Reactant); RACT (Reactant or reagent)

(benzoylation of)

IT 98-88-4, Benzoyl chloride

RL: RCT (Reactant); RACT (Reactant or reagent)

(benzoylation of methylbenzimidazole by)

IT 67264-61-3P 167281-71-2P 167281-72-3P 167281-73-4P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT

(Reactant or reagent)

(benzoylation of methylbenzimidazole by benzoyl chloride)

IT 74440-30-5P

RL: SPN (Synthetic preparation); PREP (Preparation)

(benzoylation of methylbenzimidazole by benzoyl chloride)

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

^{&#}x27;Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Emperor must precede plejseial interpretation of the year of the reign of the Year of the year of the reign of the Year of the year of the reign of the year of the reign of the year of year of the year of y

MODIFIED PTO/SB/08 A & B (08-0
Bulletin de la Societe Chimique de France (1974), (3-4, Pt. 2), 525-8
ABS Benzimidazole series. V. Behavior of 2-methylene-1,3-dimethylbenimidazoline. Alkylation and acylation reaction.
The title compound (I, X = CH2) underwent substitution with halides to give I [X = CHMe, CHCHMe2, CHCH2Ph, CHC6H3(NO2)2-2,4, CHI, 4,6-dichloro-1,3,5- triazin-2-ylmethylene, CHAc, CHBz,
CHSO2Me, CAc2, CBz2, C(SO2Me)2], some of which were isolated as the 2-alkylbenzimidazolium salts. Dimeric acylation products were obtained with ClCO(CH2)nCOCl (n = 0,2).
Me C-Ph N C-C-Ph Me
 • ні
Journal fuer Praktische Chemie (Leipzig) (1979), 321(2), 320-2
Collection of Czechoslovak Chemical Communications (1978), 43(3), 739-45
Horumon to Rinsyo (Hormones and Clinical), 46, 46-57 (1998) ABS
Gonadotropin releasing hormone is known as a hormone which controls secretion of sex hormones at the highest position, and controls secretion of anterior pituitary hormones luteinizing hormone and follicle-stimulating hormone and sex hormones in sex glands, via a receptor which is present in the anterior pituitary. Since antagonists specific and selective for this GnRH receptor regulate the action of GnRH and control secretion of subordinate LH and FSH and sex hormones, they are expected as preventive or therapeutic agents for sex hormone dependent diseases.
Molecular Endocrinology 14 671-681 2000 Identification of Phe ³¹³ of the Gonadotropin-Releasing Hormone (GnRH) Receptor as a Site Critical for the Binding of Nonpeptide GnRH Antagonists
Molecullar and Cellular Endocrin. 144 11-19 1998 Functional analysis of GnRH receptor ligand binding using biotinylated GnRH derivatives
The Prostate 20 297-310 1992 Effect of microcapsules of luteinizing hormone-releasing hormone antagonist SB-75 and somatostatin analog RC-160 on endocrine status and tumor growth in the Dunning R-3327H rat prostate cancer model.
Endocrinology 137 3430-3436 1996 Chronic administration of the luteinizing hormone-releasing hormone (LHRH) antagonist cetrorelix decreases gonadotrope responsiveness and pituitary LHRH receptor messenger ribonucleic acid levels in rats
J. Med. Chem. 2005, 48, 1169-1178
3-[(2R)-Amino-2-phenylethyl]-1-(2,6-difluorobenzyl)-5-(2-fluoro-3-methoxyphenyl)- 6-methylpyrimidin-2,4-dione (NBI 42902) as a Potent and Orally Active Antagonist of the Human Gonadotropin-Releasing Hormone Receptor. Design, Synthesis, and in Vitro and in Vivo Characterization
Bioorg. Med. Chem. Lett. 14(9) 2269-2274 2004 Synthesis and structure–activity relationships of (R)-1-alkyl-3-[2-(2-amino)phenethyl]-5-(2-fluorophenyl)-6-methyluracils as human GnRH receptor antagonists
Bioorg. Med. Chem. Lett. 15(10) 2519-2522 2005 Uracils as potent antagonists of the human gonadotropin-releasing hormone receptor without atropisomers
Curr.Opin. Drug Discovery Dev. 7, 832-847, 2004 Synthesis of orally active small-molecule gonadotropin-relea sing hormone antagonists
Bioorg. Med. Chem. Lett. 15(5) 1407-1411 2005 Efficient synthesis of bicyclic oxazolino- and thiazolino[3,2-c]pyrimidine-5,7-diones and its application to the synthesis of GnRH antagonists

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial number of the patent document, by the appropriate cyrubole as indicated on the document under WIPO Standard ST. 6 if possible. ⁵ Applicant is to indicate the Employer Handland Employer Canada ST. 6 if possible. ⁵ /V.R.G. [Page 6 of 10]

Biology Med. Chem. Lett. 15(9) 2265-2269 2005 Benzimidazoles as non-peptide luteinizing hormone-releasing hormone (LHRH) antagonists. Part 3: Discovery of 1-(1H-benzimidazol-5-yl)-3-ter-butylurea derivatives J. Med. Chem. 2004, 47, 3483-3486 3-(2-Aminostky)-1-(2-6-diflorochenzyl)-5- (2-fluoro-3-methoxyphenyl)-6-methyl-uracils as Orally Bioavailable Antagonists of the Human Gonadotropin Releasing Hormone Receptor Biorogy Med. Chem. Lett. 14 1799-1798 2004 Syntheses and structure-activity relationship studies of piperidine-substituted quinolones as nonpeptide gonadotropin releasing hormone antagonists: Biorogy Med. Chem. Lett. 14 1599-1602 2004 Elimination of antibacterial activities of non-peptide luteinizing hormone-releasing hormone (LHRH) antagonists derived from erythromycin. A J. Med. Chem. 2004, 47, 1259-1271 Synthesis and Structure-activity Relationships of 1-Arylmethyl-5-aryl-6-methyluracils as Potent Gonadotropin-teleasing Hormone Receptor Antagonists Biolong Med. Chem. Lett. 13 3317-3322 2003 Synthesis and structure-activity relationships of leap-linethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-methyluracils as antagonists of the human GnRH Receptor Biolong Med. Chem. Lett. 13 3617-3622 2003 Synthesis and structure-activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J. Med. Chem. 2004, 47, 1088-1097 Non-peptide Luteimizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis and Structure-activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists including cancers of the prostate, breast, and ovary, endometriesis, and lertifly disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3313 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and Structure-Activity relationships of novel imidazolo[1,2-a]pyr	MODIFIED PTO/SB/08 A & B (08-03
3-(2-Aminoalky)1-(2-6-diffuorobenzyl)-5- (2-fuoro-3-methoxyphenyl)-6-methyl- uracils as Orally Bioavailable Antagonists of the Human Gonadotropin Releasing Hormone Receptor Bioorg. Med. Chem. Lett. 14 1795-1798 2004 Symbeses and structure-activity relationship studies of piperidine-substituted quinolones as nonpeptide gonadotropin releasing hormone antigonists. Bioorg. Med. Chem. Lett. 14 1799-1602 2004 Elimination of antibacterial activities of non-peptide luteinizing hormone-releasing hormone (LHRH) antagonists derived from crythromycin A J. Med. Chem. 2004, 47, 1259-1271 Symthesis and Structure-Activity Relationships of 1-Arylmethyl-5-aryl-6-methyluracils as Potent Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 13 3317-3322 2003 Synthesis and Structure-activity relationships of 1-Arylmethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-methyluracils as antagonists of the human GnRH Receptor Bioorg. Med. Chem. Lett. 13 3617-3622 2003 Synthesis and structure-Activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J. Med. Chem. 2004, 47, 1085-1097 Nonepetide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Symthesis, and Biological Activity of Cladinose Replacement Antalogues J. Pharmaco. Experi. Ther. 336 588-695 2003 Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endoeri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracits as potent GnRH receptor antagonists J. Med. Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracit as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Characterization of mono- and diaminopyrimi	Benzimidazoles as non-peptide luteinizing hormone-releasing hormone (LHRH) antagonists. Part 3:
Symbess and structure—activity relationship studies of piperidine-substituted quinolones as nonpeptide gonadorropin releasing hormone antagonists: Bioorg. Med. Chem. Lett. 14 1599-1602 2004 Ellimination of antibacterial activities of non-peptide luteinizing hormone-releasing hormone (LHRH) antagonists derived from erythromycin A J. Med. Chem. 2004, 47, 1259-1271 Symbesis and Structure-Activity Relationships of 1-Arylmethyl-5-aryl-6-methyluracils as Potent Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 13 3317-3322 2003 Symbesis and Structure-Activity relationships of 1-arylmethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-methyluracils as antagonists of the human GnRH Receptor Bioorg. Med. Chem. Lett. 13 3 1617-3622 2003 Symbesis and structure-Activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J.Med. Chem. 2004, 47, 1085-1097 Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Symbesis, and Biological Activity of Cladinose Replacement Analogues J. Pharmaco. Experi. Ther. 305 688-695 2003 Gonadotropin-releasing homone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Symbesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent CnRH receptor antagonists J.Med. Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure-Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpetide luteinizing hormone-releasing hormone (LHRH) antagonists: design and	3-(2-Aminoalkyl)-1-(2,6-difluorobenzyl)-5- (2-fluoro-3-methoxyphenyl)-6-methyl- uracils as Orally
Elimination of antibacterial activities of non-peptide luteinizing hormone-releasing hormone (LHRH) antagonists derived from crythromycin A J. Med. Chem. 2004, 47, 1259-1271 Synthesis and Structure-Activity Relationships of 1-Arylmethyl-5-aryl-6-methyluracils as Potent Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 13 3317-3322 2003 Synthesis and Structure-activity relationships of 1-arylmethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-methyluracils as antagonists of the human GnRH Receptor Bioorg. Med. Chem. Lett. 13 3617-3622 2003 Synthesis and Structure-Activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J.Med. Chem. 2004, 47, 1085-1097 Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis, and Biological Activity of Cladinose Replacement Analogues J. Pharmaco. Experi. Ther. 305 688-695 2003 Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and lertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med. Chem. 2003, 46, 2003-20026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and sinuture-Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2073-2072 2002 A new class of potent nonpeptide luteriinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylmidiazol[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-7?? Bioorg. Med. Chem. Lett. 12 3633-3639 2002 Characterization of nono and diami	Syntheses and structure-activity relationship studies of piperidine-substituted quinolones as nonpeptide
Synthesis and Structure-Activity Relationships of 1-Arylmethyl-5-aryl-6-methyluracils as Potent Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg, Med. Chem. Lett. 13 3317-3322 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-methyluracils as antagonists of the human GnRH Receptor Bioorg, Med. Chem. Lett. 13 3617-3622 2003 Synthesis and structure-Activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J. Med. Chem. 2004, 47, 1085-1097 Nonpeptide Luterinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis, and Biological Activity of Cladinose Replacement Analogues J. Pharmaco. Experi. Ther. 305 688-695 2003 Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J. Med. Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure-Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2183-2187 2002 Design, synthesis and structure-Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J. Med. Chem. Lett. 12 2183-2187 2002 Characterization of mono- and diaminopyrimidine-ones Drugs of the Future 2003, 28, 121-727 Bioorg. Med. Chem. Lett. 12 3633-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antago	Elimination of antibacterial activities of non-peptide luteinizing hormone-releasing hormone (LHRH)
Synthesis and Structure-activity relationships of 1-arylmethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-methyluracils as antagonists of the human GnRH Receptor Bioorg, Med. Chem. Lett. 13 3617-3622 2003 Synthesis and structure-Activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J.Med.Chem. 2004, 47, 1085-1097 Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis, and Biological Activity of Cladinose Replacement Analogues J. Pharmaco. Experi. Ther. 305 688-695 2003 Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure-Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure-Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3353-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists	Synthesis and Structure-Activity Relationships of 1-Arylmethyl-5-aryl-6-methyluracils as Potent
Synthesis and structure—Activity relationships of thieno[2,3-a/]pyrimidine-2,4-dione derivatives as potent GnRH receptor antagonists J.Med.Chem. 2004, 47, 1085-1097 Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis, and Biological Activity of Cladinose Replacement Analogues J. Pharmaco. Experi. Ther. 305 688-695 2003 Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure—Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med Chem. 2003, 46, 2023-2026 Identification of 1-arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure—Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonepetide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Synthesis and Structure–activity relationships of 1-arylmethyl-3-(1-methyl-2-amino)ethyl-5-aryl-6-
Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design, Synthesis, and Biological Activity of Cladinose Replacement Analogues J. Pharmaco. Experi. Ther. 305 688-695 2003 Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure-Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure-Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure-Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-2?? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 329-3332 2002	Synthesis and structure—Activity relationships of thieno[2,3-d]pyrimidine-2,4-dione derivatives as potent
Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and fertility disorders J. Clin. Endocri. Metab. 88 1697-1704 2003 Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure-Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure-Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure-Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3335-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Nonpeptide Luteinizing Hormone-Releasing Hormone Antagonists Derived from Erythromycin A: Design,
Bioorg. Med. Chem. Lett. 13 3311-3315 2003 Synthesis and Structure—Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure—Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Gonadotropin-releasing hormone (GnRH) receptor antagonists have potential in treating numerous hormone-dependent pathologies including cancers of the prostate, breast, and ovary, endometriosis, and
Synthesis and Structure—Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 2023-2026 Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure—Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	J. Clin. Endocri. Metab. 88 1697-1704 2003
Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone Receptor Antagonists Bioorg. Med. Chem. Lett. 12 2179-2183 2002 Synthesis and initial structure—Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Synthesis and Structure–Activity relationships of 1-arylmethyl-3-(2-aminopropyl)-5-aryl-6-methyluracils as
Synthesis and initial structure—Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists Bioorg. Med. Chem. Lett. 12 2185-2187 2002 Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Identification of 1-Arylmethyl-3- (2-aminoethyl)-5-aryluracil as Novel Gonadotropin-Releasing Hormone
Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent GnRH receptor antagonists J.Med.Chem. 2003, 46, 113-124 Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Synthesis and initial structure–Activity relationships of a novel series of imidazolo[1,2-a]pyrimid-5-ones as
Bioorg. Med. Chem. Lett. 12 2073-2077 2002 A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	Design, synthesis and structure—Activity relationships of novel imidazolo[1,2-a]pyrimid-5-ones as potent
A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and synthesis of 2-phenylimidazo[1,2-a]pyrimidin-5-ones Drugs of the Future 2003, 28, 121-??? Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	J.Med.Chem. 2003, 46, 113-124
Bioorg. Med. Chem. Lett. 12 3635-3639 2002 Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	A new class of potent nonpeptide luteinizing hormone-releasing hormone (LHRH) antagonists: design and
Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing hormone (GnRH) receptor antagonists Bioorg. Med. Chem. Lett. 12 3329-3332 2002	
	Characterization of mono- and diaminopyrimidine derivatives as novel, nonpeptide gonadotropin releasing
Modification of the pyridine moiety of non-peptidyl indole GnRH receptor antagonists	Bioorg. Med. Chem. Lett. 12 3329-3332 2002 Modification of the pyridine moiety of non-peptidyl indole GnRH receptor antagonists

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

^{&#}x27;Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede Adsertial Implication of the year of the reign of the Applicant is to indicate here if English language Translation is attached.

[Decay 7 = 6.10]

	MODIFIED PTO/SB/08 A &	B (08-0
	Bioorg. Med. Chem. Lett. 12 3467-3470 2002	
	The discovery of novel small molecule non-peptide gonadotropin releasing hormone (GnRH) receptor	
	antagonists	
	Bioorg. Med. Chem. Lett. 12 3491-3495 2002	
	A novel synthesis of 7-aryl-8-fluoro-pyrrolo[1,2-a]pyrimid-4-ones as potent, stable GnRH receptor	
	antagonists	
	J.Med.Chem. 2001, 44, 917-922	
	A Potent, Nonpeptidyl 1 <i>H</i> -Quinolone Antagonist for the Gonadotropin-Releasing Hormone Receptor	
	Bioorg. Med. Chem. Lett. 12 399-402 2002	
ļ	Initial Structure—Activity Relationship Studies of a Novel Series of Pyrrolo[1,2-a]pyrimid-7-ones as GnRH	
	Receptor Antagonists	
	Bioorg. Med. Chem. Lett. 12 403-406 2002	
	A Novel Synthesis of 2-Arylpyrrolo[1,2-a]pyrimid-7-ones and Their Structure–Activity Relationships as	
	Potent GnRH Receptor Antagonists	
.	Tetraherdron Lett. 42 6441-6445 2001	
	Total syntheses of 6- and 7-azaindole derived GnRH antagonists	_
	Tetraherdron Lett. 42 6459-6461 2001	
	Synthesis of chiral β-methyl tryptamine-derived GnRH antagonists	
	Bioorg. Med. Chem. Lett. 11 2597-2602 2001 Orally bioavailable, indole-based nonpeptide GnRH receptor	
	antagonists with high potency and functional activity	
1	Bioorg. Med. Chem. Lett. 11 1723-1726 2001	
	Substituted Indole-5-carboxamides and -acetamides as Potent Nonpeptide GnRH Receptor Antagonists	
<u></u>	Bioorg. Mcd. Chem. Lett. 11 1727-1731 2001	-
	Potent nonpeptide GnRH receptor antagonists derived from substituted indole-5-carboxamides and -	
	acetamides bearing a pyridine side-chain terminus	
	Tetraherdron 57 5233-5241 2001	
	A convergent synthesis of (S)-β-methyl-2-aryltryptamine based gonadotropin releasing hormone	
	antagonists	
<u> </u>		
	Bioorg. Med. Chem. Lett. 15 2519-2522 2005	
	Uracils as potent antagonists of the human gonadotropin-releasing hormone receptor without atropisomers	
	Bioorg. Med. Chem. Lett. 15 4363-4366 2005	
	Synthesis of aryl-1,2,4-triazine-3,5-diones as antagonists of the gonadotropin-releasing hormone receptor	
	Bioorg, Med. Chem. Lett. 15 3685-3690 2005	
1	Structure–activity relationships of 1,3,5-triazine-2,4,6-triones as human gonadotropin-releasing hormone	
	receptor antagonists	
	Bioorg. Med. Chem. Lett. 15 799-803 2005	
	Benzimidazole derivatives as novel nonpeptide luteinizing hormone-releasing hormone (LHRH)	
	antagonists. Part 1: Benzimidazole-5-sulfonamides c	
	Bioorg, Med. Chem. Lett. 15 805-807 2005	
	Benzimidazole derivatives as novel nonpeptide luteinizing hormone-releasing hormone (LHRH)	
	antagonists. Part 2: Benzimidazole-5-sulfonamides	
	Bioorg. Med. Chem. Lett. 14 2269-2274 2004	
	Synthesis and structure–activity relationships of (R)-1-alkyl-3-[2-(2-amino)phenethyl]-5-(2-fluorophenyl)-	
	6-methyluracils as human GnRH receptor antagonists	
	Bioorg. Med. Chem. Lett. 14 5599-5603 2004	
	Identification of neutral 4-O-alkyl quinolone nonpeptide GnRH receptor antagonists	
	Bioorg. Med. Chem. Lett. 9 2615-2620 1999	
	Identification and Initial Structure-Activity Relationships of a Novel Non-Peptide Quinolone GnRH	
	Receptor Antagonist	
	Bioorg. Med. Chem. Lett. 9 2621-2624 1999	
	Investigation of the 4-O-Alkylamine Substituent of Non-Peptide Quinolone GnRH Receptor Antagonists A	

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede the serial impropriate patent patents of the patent of the patents of the p

	Modified Flore	B/08 A & B (08-
	Bioorg. Med. Chem. Lett. 10 1723-1727 2000	
	Annual Reports in Medicinal Chemistry vol.39 99-110 (2004)	
	J.Med.Chem. 1998, 41, 4190-4195	
j	Discovery of a Novel, Potent, and Orally Active Nonpeptide Antagonist of the Human Luteinizing	
	Hormone-Releasing Hormone (LHRH) Receptor	
	Angew. Chem. Int. Ed. Engl. 1997 36 2148-2161	
	Chemistry and Molecular Biology in the Search for New LHRH Antagonists	
	J.Med.Chem. 1989, 32, 2036-2038	
	Zeitschrift fuer Chemie (1980), 20(3), 96-7	ļ
	Synthesis and reactions of 2,2-diacylketene heteroacetals	
	ABS	
	H Me Ph	
	(PhCO) 2C X	
	I NC S CN II	
	H I NC S S CN II	
	Bz Bz Bz	
1	Bz Ph	
	MeS SMe III MeS Ph	
	Mes See III Fil IV	
	The reactions of thioacetals and dithiolates were described. Thus, (RCO)2C:C(SMe)2 (R = Me, Ph)	
	reacted with PhCH2NH2 to give (RCO)2C:CR1R2 (R1 = SMe, R2 = NHCH2Ph; R1 = R2 = NHCH2Ph)	
	and (PhCO)2C:C(SMe)2 reacted with dinucleophiles, e.g., (H2NCH2)2 and o-(H2N)2C6H4, to give	
	cyclic heteroacetals, e.g. I (X = CH2CH2 o-C6H4). Alkylation of MeCO(PhCO)C:C(S-)2 with ClCH2CN	
	gave thienothiophene II via an open-chain S,S-acetal and subsequent cyclization. PhCOCH2COCH2Ph	
	reacted with CS2 and NaH to cleave Na2S; alkylation of the product with Mel gave thiopyranone III.	
	Treating (PhCOCH2)2 with PhNCS gave PhCOC(CH2COPh):C(NHPh)S- which was cyclized and	
	methylated to give pyrrole IV.	ļ
	The Prostate(1992) 20 297-310	
	Bioorg. Med. Chem. Lett. 11 515-517 2001	
	Bioorg. Med. Chem. Lett. 11 509-513 2001	
	Bioorg. Med. Chem. Lett. 12 93-96 2002	
	Bioorg. Med. Chem. Lett. 12 827-832 2002	
	J.Med.Chem.(2006), 49, 3809-3825	
	Drug of the Future(1999) 24(4) 393-403	
	J. Peptide Res.(2005)65 284-291	
	Int. J. Peptide Protein Res.(1988) 32 425-435	
	J. Clinical Endocrinology and Metabolism(1997) 82 5 1403-1408	
	J. Clinical Endocrinology and Metabolism(1992) 75 2 393-398	
	Current Pharma. Design(2003) 9 381-390	
	Mol. Endocrinology(2000) 14(7) 1099-1115	
	Bioorg. Med. Chem. Lett. 12 827-832 2002	
	JMC(2001) 44 453-467	
	EUR J MED CHEM(1997) 32 927-940	· ·
	JMC(2000) 43 2831-2836	
	Bioorganic & Medicinal Chemistry Letters(2005) 15 1609-1612	
		
	PNAS(2002) 99 961-965	
	JMC(1997) 40 3739-3748	
	JMC(2000) 43 784-796	
	JMC(2000) 43 797-806 JMC(2000) 43 807-818	

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

¹Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede and squal number of the patent programment by the appropriate symbols as indicated on helder under WIPO Standard ST. (1) Possible 1. /V.R.G. Applicant is to indicate here if English language Translation is attached.

JMC(2000) 43 819-828	
JMC(2002) 45 1026-1034	
JMC(2005) 48 4851-4860	
J. Endocrinol., Feb 1988; 116: 241 - 246.	
Endocrine Rev.(1986) 7 1 44-66	
Current Sci.(1987) 56 7 325-327	
J Endocrinol 1986 108: 101-107	
Neuroendocrinology(1985) 40 246-252	
Endocrinology 1983 113: 195-199	
Biology of Reproduction(1999) 61 1468-1479	
Mol. Endocrinology(1997) 11 11 1659-1668	
Biology of Reproduction(1995) 53 724-731	
Comp. Biochem. Physiol.(1994) 108C 1 129-135	
Current Opinion in Endocrinology & Diabetes(2000) 7 350-356	
J Reprod. Fert.(1992) 96 865-874	
J Clin Endocrinol Metab 1992 75: 1220-1225	
TEM(1992) 3 7 259-263	
J Clin Endocrinol Metab 1994 78: 121-125	
Hormone Res.(1987) 28 88-103	
andrologia(1990) 22 567-573	
Neuroendocrinology(1989) 50 158-16421	
Neuroendocrinology(1989) 50 158-16421	

	Malayla Dadvievan appela/		12/07/2000
Examiner Signature	/Valerie Rodriguez-garcia/	Date Considered	12/0//2009

^{*}EXAMINER: Initial if reference considered, whether or not citation is in conformance with MPEP 609. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.

Applicant's unique citation designation number (optional). ²See Kind Codes of USPTO Patent Documents at www.uspto.gov, MPEP 901.04 or follow the hyperlink from the title of the document to the intranet. ³ Enter Office that issued the document, by the two-letter code (WIPO Standard ST. 3). ⁴For Japanese patent documents, the indication of the year of the reign of the Emperor must precede Acts drial paths for the part of the reign of the Emperor must precede Acts drial paths for the part of the reign of the Emperor must precede Acts drial paths for the path of the path of the paths for the path of the path of the paths for the path of the pat